

ROMAX

Contains Coumatetralyl 0.0375% w/w

DIRECTIONS FOR USE

Place bait where there are signs of rat activity, like fresh droppings, inside or near burrows, runways or feeding places. Place bait throughout the infested area.



For rat use 1-2 soft blocks (sachets) per baiting point depending upon the severity of the infestation.

Make frequent inspections of the bait points. Replace any eaten bait. If all bait has been eaten from certain areas, increase the quantity of bait by placing more bait points. Do not increase the bait point size. The resistance status of the population should be taken into account when considering the choice of rodenticide to be used.

Protect bait points from non-target animals and from water, preferably using commercially available bait stations. Where these are not available place the bait inside pieces of drainage pipe or under sections of slate, board or corrugated iron, suitably weighted. Soft block may be fixed by 'spiking' onto bars or wiring inside bait stations. Make frequent inspections of the bait points during the first 10-14 days and replace any bait that has been eaten by rodents or has been damaged by water or contaminated by dirt.

GB79413580A - ARTICLE 79288107



5 015861 985101

Romax Rat CP contains the bittering agent Bitrex™, the human taste deterrent, providing greater bait safety without loss of effectiveness.

The bait packaging material is biodegradable.

Romax is a registered trademark of JV Barrett and Co.

Barrettine**Environmental Health**

St. Ivel Way, Warmley, Bristol BS30 8TY
Tel: 0117 967 2222 Fax: 0117 961 4122
email: beh@barrettine.co.uk

STATUTORY CONDITIONS RELATING TO USE.
Application rate: Please see directions for use.**FOR USE ONLY BY PROFESSIONAL OPERATORS**

The (COSHH) Control of Substances Hazardous to Health Regulations 2002 may apply to the use of this product at work.

AVOID ALL CONTACT BY MOUTH.

PREVENT ACCESS TO THIS PRODUCT by children, birds and non-target animals [particularly dogs, cats, pigs, poultry].

Search for and remove rodent bodies at frequent intervals*. Collect and dispose of the remains of bait and any remaining rodent bodies after treatment (unless used in sewers).

You must ensure that you comply with legislation regarding the correct disposal of waste. For further guidance, contact the Environment Agency or your local Scottish Environment Protection Agency (SEPA) office.

HARMFUL TO WILDLIFE.

DO NOT PLACE THIS PRODUCT where food, feed or water could become contaminated (except when used in sewers).

(* Intervals will vary depending on formulation and treatment regime; therefore the user should refer to and comply with the manufacturer's guidelines).

READ ALL PRECAUTIONS BEFORE USE.

HSE No. ?????

PRECAUTIONS

Wild mammals may be at greater risk if this product is not used in accordance with the label. **IF YOU FEEL UNWELL**, seek medical advice (show the label where possible). **WASH HANDS AND EXPOSED SKIN** before meals and after use. Keep only in the original container. Keep container tightly closed. **KEEP IN SAFE PLACE EMPTY CONTAINER COMPLETELY** and dispose of safely. **STORE UNUSED SACHETS** in a safe place. **DO NOT STORE** part used sachets. Where appropriate, secure the bait sachet at its placement site to reduce the likelihood of its removal by rodents [or by surges of water if intended for use in sewers]. **COUMATETRALYL** is an indirect anticoagulant. Phytomenadione, Vitamin K₁ is antidotal. Determine prothrombin times not less than eighteen hours after consumption. If elevated, administer Vitamin K₁ until prothrombin time normalises. Continue determination of prothrombin time for two weeks after withdrawal of antidote and resume treatment if elevation occurs in that time.

This product is approved under The Control of Pesticides Regulations 1986 (as amended) for use as directed. Use only in accordance with the directions on the label.