

READ THE LABEL BEFORE USE. Using this product in a manner that is inconsistent with the label may be an offence. Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.

Hazard and precautionary statements:

WARNING



GHS08

H373: May cause damage to organs through prolonged or repeated exposure.

Precautionary Statements

- P102 - Keep out of reach of children
- P270 - Do not eat, drink or smoke when using this product
- P273 - Avoid release to the environment
- P301 + 312 - If swallowed: Call a poison center or doctor/physician if you feel unwell
- P501 - Dispose of contents/container in accordance with national regulations

Directions for use:

For professional use only indoors and outdoors around buildings, open areas, and in waste dumps. Prevent access to bait by children, birds and non-target animals (particularly dogs, cats, pigs and poultry). Determine areas where rats and mice will likely find and consume bait. Generally, these areas are along walls, by gnawed openings, in corners, and concealed places.

Rats:

Place one to five place packs in a bait station or covered bait point. Place at 5 to 10 metre intervals per placement where rats or their signs have been observed. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

Mice:

Place one place pack in a bait station or covered bait point. Place at 5 to 10 metre intervals per placement where mice or their signs have been observed. Maintain an uninterrupted supply of fresh bait for 15 days or until signs of mice activity cease.

Public Use Areas: For products to be used in public areas the following safety precaution shall be carried on the label, packaging, or accompanying leaflet: When Contrac Rodenticide is used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant as well as indicating the first measures to be taken in case of poisoning must be made available alongside the baits. When tamper resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed. Antidote vitamin K₁ (under medical supervision), UK medical professional should contact the National Poisons Information Service (www.npis.org) for further advice.

Additional Use Instructions: Wash hands and exposed skin before meals and after use. Contrac Rodenticide may be applied by professional users only when appropriate chemical resistant gloves (EN374) are used. Search for and remove dead rodents a frequent intervals during treatment, at least as often as when baits are checked and/or replenished. Daily inspections may be required in some circumstances. Prevent access to bait by children, birds and non-target animals (particularly dogs, cats, pigs and poultry). Baits must be securely deposited in a way so as to minimise the risk of consumption by other animals or children. Where possible, secure baits so that they cannot be dragged away.

 Bell Laboratories Supports



Campaign for Responsible Rodenticide Use

6562-1

CONTRAC[®]

RODENTICIDE

KILLS RATS & MICE

A ready to use pelleted bait containing Bromadiolone for use as a rodenticide by professionals for the control of rats and mice indoors and outdoors around buildings, open areas, and waste dumps for the protection of stored products, food, health, and material protection (e.g. historical buildings, technical objects, etc.)

TO BE USED ONLY BY PROFESSIONAL USERS HOLDING CERTIFICATION
DEMONSTRATING COMPLIANCE WITH UK RODENTICIDE
STEWARDSHIP REGIME REQUIREMENTS

NOT for Amateur Sale

Active Substance: Bromadiolone 0.05 g/kg (0.005% w/w)
(CAS No. 28772-56-7)

Contains Denatonium Benzoate – bittering agent
Use Biocides Safely and Sustainably

Authorisation No.: UK-2015-0919

NET WEIGHT: 5 kg
100 x 50g place packs

Authorisation holder:

Bell Laboratories, Inc. – European Division
Chaucer House, Chaucer Road
Sudbury, Suffolk
CO10 1LN ENGLAND
Tel: +44 1787379295
Email: emea@belllabs.com

Manufactured By:

 **Bell**
LABORATORIES, INC.
www.belllabs.com
Madison, WI 53704 U.S.A.
MADE IN USA

APPROVED

911010

Resistance and Long Term Use:

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients. Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits. In most cases, anticoagulant bait should have achieved control within 35 days. Should activity continue beyond this time, the likely cause should be determined.

First Aid and Safety information

In case of accident or if taken unwell, seek medical advice immediately (show the label if possible). Bromadiolone is an anticoagulant which may produce bleeding; this may be delayed for several days after exposure. If there is no active bleeding the INR (prothrombin time) should be measured on presentation and 48–72 hours after exposure. If the INR is greater than 4, administer Vitamin K₁ (phytomenadione) 5–10 mg by slow intravenous injection (100 µg/kg body weight for a child). Treatment with phytomenadione (orally or intravenously) may be required for several weeks. The advice of the National Poisons Information Service (<http://www.npis.org>) should be sought, particularly if active bleeding occurs.

Antidote – Vitamin K₁ (under medical supervision). UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.

In case of contact with skin, wash with soap and water. Remove and launder any contaminated clothing.

In case of contact with eyes, remove contact lenses, if present, and rinse the eye slowly and gently with water for 15–20 minutes. Seek medical advice immediately.

In case of ingestion or if swallowed, seek medical advice immediately.

In case of pet emergency, contact your veterinarian immediately.

Storage Instructions:

Store in the original container in a cool, dry, well ventilated place. Do not reuse the container for any other purpose. Shelf life up to 2 years.


Directions for safe disposal:

Remove all baits after treatment and dispose of them in accordance with local requirements. Dead rodents and bait found outside bait stations should be disposed of using protective gloves. Dispose of dead rodents in accordance with local requirements. Poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (<http://www.environment-agency.gov.uk>) or SEPA (<http://www.sepa.org.uk>).

Batch Number, Date of Manufacture, and Expiry Date are printed on the packaging.

When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements.

052716/05-16



3 00 30835 50136 7