



You know when
you're in safe hands!

www.killgerm.com

Bird Free[®] passes with flying colours!



[®] 小島 **BIRD FREE** 

Protecting the
urban environment

Killgerm Chemicals Ltd., Wakefield Road, Ossett,
West Yorkshire, WF5 9AJ, UK
t +44 (0) 1924 268420 e sales@killgerm.com



BIRD FREE[®] IS BACK!!

- Ready to use dishes
- Quick, easy, mess-free installation
- Making working at heights safer and easier
- Firm texture enables use on pitched roofs and angled surfaces
- Discreet low profile dishes
- Keeps pigeons away from structures without harming them
- **NOW WITH HSE AUTHORISATION**



Killgerm®

You know when
you're in safe hands!

www.killgerm.com

INSTALLATION MANUAL

® 小島 **BIRD FREE**

Protecting the urban environment

Directions for Use

Bird Free® may be used to deter feral pigeons from roosting or nesting on buildings and other structures. Bird Free® is supplied in ready-to-use plastic dishes (65mm diameter; 8mm high). These should be affixed to surfaces with commercially available silicone adhesive (e.g. Avisil / Silirub) after removing covers. Allow surface to dry before applying silicone adhesive. Magnetic dishes are also available.



Installation guidelines for feral pigeons

1. Clean site thoroughly using:

- **PX-Ornikill** (See label for dilution rates)

» for disinfecting bird droppings


- **Surface Cleaner ADH003-2**

» for cleaning surfaces prior to installation


(See "Cleaning the site prior to application of Bird Free®", right)

2. Fix dishes as shown below.


High pressure/nesting sites: 15cm (centre to centre)



Medium pressure/night roosts: 20cm (centre to centre)



Low pressure/day roosts: 25cm (centre to centre)



Cleaning the site prior to application of Bird Free®

It is essential to clean the application site of all nests, faeces and other organic debris before Bird Free® is applied. (Failure to do so will compromise efficacy.)

Routine cleaning of droppings where there are no nests or major accumulation of faeces (e.g. from window sills) can be done with scrapers and wire brushes. Simple precautions to reduce direct contact with droppings, such as wearing disposable gloves and disposable coveralls and other suitable Personal Protective Equipment should be taken.

When cleaning long-standing nesting or night roosting sites, where there is a major accumulation of faeces, full PPE, including masks, should be worn. Precautions should be taken to ensure that members of the public are not exposed to airborne dust or other debris. Surfaces should be wetted with biocide, e.g. PX-Ornikill, or water mixed with detergent where faeces are present to limit generation of airborne dust and aerosol mists during cleaning. If a pressure washer is used, dust and aerosol control measures, such as containing the area with plastic sheeting, should be taken. Alternatively the wetted faeces and debris can be removed with a shovel. The wet material should be collected in heavy-duty plastic bags, or some other type of secure container, and suitable arrangements made for disposal.

Wash skin and clothing after cleaning bird debris and before eating or drinking. Do not touch or rub eyes while or after handling Bird Free®. When cleaning large amounts of bird debris, operatives should be informed of the possible health risks involved. Persons with compromised immune systems should avoid the cleaning operation.

For technical support or other questions, contact:

Killgerm Chemicals Ltd., Wakefield Road, Ossett, West Yorkshire, WF5 9AJ, UK

t +44 (0) 1924 268420 **e** sales@killgerm.com

